

Health Indicator Sortable Stats About the Data

Heart Disease Death Rate (2007)

Indicator: Age adjusted rate of heart disease deaths per 100,000 population

Source: NCHS National Vital Statistics System (NVSS) <http://www.cdc.gov/nchs/fastats/heart.htm>

About this Indicator: Rates were calculated per 100,000 population. Age-adjustments were calculated using the 2000 population standard. NVSS deaths from diseases of heart, codes include (I00-I09, I11, I13, and I20-I51)

Additional Resources and Links:

MMWR: Coronary Heart Disease and Stroke Deaths --- United States, 2006

http://www.cdc.gov/mmwr/preview/mmwrhtml/su6001a13.htm?s_cid=su6001a13_w

Health Indicators Warehouse: Heart disease deaths

http://www.healthindicators.gov/Indicators/Heart-disease-deaths_83/Profile

Quick Health Data Online: (May contain city and/or county level data)

http://www.healthstatus2010.com/owh/select_variables.aspx

Stroke Death Rate (2007)

Indicator: Age adjusted rate of stroke deaths per 100,000 population

Source: NCHS National Vital Statistics System (NVSS) <http://www.cdc.gov/nchs/fastats/stroke.htm>

About this Indicator: Rates were calculated per 100,000 population. Age-adjustments were calculated using the 2000 population standard. NVSS Deaths from Stroke (cerebrovascular disease), codes include (ICD 10 I60-I69).

Additional Resources and Links:

MMWR: Coronary Heart Disease and Stroke Deaths --- United States, 2006

http://www.cdc.gov/mmwr/preview/mmwrhtml/su6001a13.htm?s_cid=su6001a13_w

Health Indicators Warehouse: Stroke deaths (including county level data)

http://www.healthindicators.gov/Indicators/Stroke-deaths_881/Profile

Quick Health Data Online: (May contain city and/or county level data)

http://www.healthstatus2010.com/owh/select_variables.aspx

HIV Diagnosis Rate (2009)

Indicator: Rate of persons diagnosed with HIV infection per 100,000 population

Source: CDC 2009 HIV Surveillance Report

<http://www.cdc.gov/hiv/surveillance/resources/reports/2009report/index.htm#technical>

About this Indicator: This figure does not include persons living with HIV who have NOT been diagnosed (20% of all persons living with HIV). These data use the 2008 revised HIV case definition to classify HIV infection which increases the specificity of the case definition by requiring laboratory-confirmed evidence of HIV infection. As of December 2008 HIV cases are reported for all 50 states, DC and 5 territories that require confidential reporting.

Additional Resources and Links:

CDC Vital Signs: HIV Testing in the US

<http://www.cdc.gov/vitalsigns/HIVTesting/>

MMWR: HIV Testing and Diagnosis Among Adults --- United States, 2001--2009

http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5947a3.htm?s_cid=mm5947a3_w



Health Indicator Sortable Stats About the Data

Central Line-Associated Bloodstream Infection- Standardized Infection Ratio (CLABSI-SIR) (2010)

Indicator: Central Line-Associated Bloodstream Infections -Standardized Infection Ratios

Source: National Healthcare Safety Network (NHSN); CDC First State-specific HAI Summary Data Report

http://www.cdc.gov/hai/pdfs/stateplans/SIR_05_25_2010.pdf

http://www.hhs.gov/ash/initiatives/hai/actionplan/hhs_hai_action_plan_final_06222009.pdf

About this Indicator: Based on Healthcare Facilities currently report using NHSN (17 states). In HAI data analysis, the SIR compares the actual number of HAIs in a facility or state with the predicted number based on the baseline U.S. experience (i.e., standard population), adjusting for several risk factors that have been found to be most associated with differences in infection rates.

Additional Resources and Links:

CDC Vital Signs: Making Health Care Safer: Reducing bloodstream Infections

<http://www.cdc.gov/vitalsigns/HAI/index.html>

Motor Vehicle Death Rate (2009)

Indicator: Fatality rate per 100,000 population

Source: Fatality Analysis Reporting System (FARS)/ NHTSA

<http://www-fars.nhtsa.dot.gov/Trends/TrendsGeneral.aspx>

About this Indicator: Based on traffic related vehicle crashes in the 50 states and DC that occur on public roadways and involves a fatality.

Additional Resources and Links:

Health Indicators Warehouse: Motor Vehicle Deaths (including county level data)

http://www.healthindicators.gov/Indicators/Motor-vehicle-deaths_18/Profile

Opioid Analgesics Deaths (2007)

Indicator: Number of fatal poisoning deaths involving Opioid Analgesics

Source: NCHS National Vital Statistics System.

<http://www.cdc.gov/nchs/data/databriefs/db22.htm>

About this Indicator: NVSS Deaths from Poisoning deaths involving Opioid analgesics, ICD-10 codes include X40-X49, X60-X69, X85-X90, Y10-Y19, Y35.2, or *U01(.6-.7) as an underlying cause of death and any of the codes T40.2-T40.4 as a multiple cause. Rates were calculated per 100,000 population. Age-adjustments were calculated using the 2000 population standard.

Additional Resources and Links:

MMWR Quick Stats: Number of Poisoning Deaths* Involving Opioid Analgesics and Other Drugs or Substances --- United States, 1999-2007

http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5932a6.htm?s_cid=mm5932a6_w

Teen Birth Rate (2009)

Indicator: Birth Rate for teens ages 15-19 per 1,000 female population (preliminary data).

Source: NCHS National Vital Statistics System. <http://www.cdc.gov/nchs/VitalStats.htm>

http://www.cdc.gov/nchs/data/nvsr/nvsr59/nvsr59_01.pdf



Health Indicator Sortable Stats About the Data

About this Indicator: Race/Ethnicity Data based on 2008 final birth rates. 2009 rates are based on preliminary data.

Additional Resources and Links:

NCHS Data Briefs: US Teen Birth Rate Resumes Decline

<http://www.cdc.gov/nchs/data/databriefs/db58.htm#Ref1>

CDC Vital Signs: Preventing Teen Pregnancy in the US

<http://www.cdc.gov/vitalsigns/TeenPregnancy/index.html>

Health Indicators Warehouse: Teen Birth Rate (including county level data)

http://www.healthindicators.gov/Indicators/Teen-birth-rate_22/Profile

Diagnosed High Cholesterol (2009)

Indicator: Percent of adults who have had their blood cholesterol checked and have been told it was high

Source: Behavioral Risk Factor Surveillance System 2009

<http://apps.nccd.cdc.gov/BRFSS/list.asp?cat=CA&yr=2009&qkey=4392&state=All>

About this Indicator: Includes all 50 states and DC. National data based on median state prevalence of 50 States and DC (median %=37.5).

Additional Resources and Links:

CDC Vital Signs: High Blood Pressure and Cholesterol

<http://www.cdc.gov/vitalsigns/CardiovascularDisease/index.html>

MMWR Prevalence, Treatment, and Control of High Levels of Low-Density Lipoprotein Cholesterol --- United States, 1999--2002 and 2005--2008

http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6004a5.htm?s_cid=mm6004a5_w

SMART: BRFSS City and County Data (2009): <http://apps.nccd.cdc.gov/BRFSS-SMART/ListMMSAQuest.asp?yr2=2009&MMSA=All&cat=CA&qkey=4392&grp=0>

Diagnosed Hypertension (2009)

Indicator: Percent of Adults (age 18+) ever told by health professional you have High blood pressure

Source: Behavioral Risk Factor Surveillance System 2009

<http://apps.nccd.cdc.gov/BRFSS/list.asp?cat=HA&yr=2009&qkey=4420&state=All>

About this Indicator: Includes all 50 states and DC. National data based on median state prevalence of 50 States and DC (median %=28.7).

Additional Resources and Links:

CDC Vital Signs: High Blood Pressure and Cholesterol

<http://www.cdc.gov/vitalsigns/CardiovascularDisease/index.html>

MMWR: Prevalence, Treatment, and Control of Hypertension --- United States, 1999--2002 and 2005--2008

http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6004a4.htm?s_cid=mm6004a4_w



Health Indicator Sortable Stats

About the Data

SMART: BRFSS City and County Data (2009): <http://apps.nccd.cdc.gov/BRFSS-SMART/ListMMSAQuest.asp?yr2=2009&MMSA=All&cat=HA&qkey=4420&grp=0>

Health Indicators Warehouse: High blood pressure (including county level data)
http://www.healthindicators.gov/Indicators/High-blood-pressure_123/National_0/Profile

Medicated Hypertension (2009)

Indicator: Percent of Adults (age 18+) currently taking medicine for high blood pressure (hypertension)

Source: Behavioral Risk Factor Surveillance System 2009

About this Indicator: Percent is based on a subset of persons having been told by health professional that they have high blood pressure. Data includes all 50 states and DC. National data based on median state prevalence of 50 States and DC (median %=78.1)

Additional Resources and Links:

CDC Vital Signs: High Blood Pressure and Cholesterol
<http://www.cdc.gov/vitalsigns/CardiovascularDisease/index.html>

MMWR: Prevalence, Treatment, and Control of Hypertension --- United States, 1999--2002 and 2005--2008
http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6004a4.htm?s_cid=mm6004a4_w

Adult Smoking (2010)

Indicator: Percent of Adults (age 18+) who currently smoke cigarette

Source: Behavioral Risk Factor Surveillance System 2010
<http://apps.nccd.cdc.gov/BRFSS/list.asp?cat=TU&yr=2010&qkey=4396&state=All>

About this Indicator: Includes all 50 states. National data based on median state prevalence of 50 States and DC (median %=17.3).

Additional Resources and Links:

CDC Vital Signs: Tobacco Use: Smoking and Secondhand Smoke
<http://www.cdc.gov/vitalsigns/TobaccoUse/Smoking/index.html>
<http://www.cdc.gov/vitalsigns/TobaccoUse/SecondhandSmoke/index.html>

MMWR: Current Cigarette Smoking Among Adults Aged ≥ 18 Years --- United States, 2009 and Nonsmokers' Exposure to Secondhand Smoke --- United States, 1999--2008
http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5935a3.htm?s_cid=mm5935a3_w
http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5935a4.htm?s_cid=mm5935a4_w

SMART: BRFSS City and County Data (2009): <http://apps.nccd.cdc.gov/BRFSS-SMART/ListMMSAQuest.asp?yr2=2009&MMSA=All&cat=TU&qkey=4396&grp=0>

Health Indicators Warehouse: Smoking (including county level data)
http://www.healthindicators.gov/Indicators/Smoking_13/Profile

Youth Smoking (2009)

Indicator: Percent of high school students who smoked cigarettes on at least 1 or more days in last 30 days

Source: Youth Risk Behavior Surveillance System (YRBSS)



Health Indicator Sortable Stats About the Data

<http://apps.nccd.cdc.gov/YouthOnline/App/Default.aspx?SID=HS>

About this Indicator: National median based on data from 42 states using YBRSS survey of adolescent's grades 9-12 (median%=19.5). Data not available in CA, DC, IA, MN, NE, OH, OR, VA, and WA.

Additional Resources and Links:

MMWR: Youth Risk Behavior Surveillance—United States, 2009

<http://www.cdc.gov/mmwr/preview/mmwrhtml/ss5905a1.htm>

Health Indicators Warehouse: Adolescent use of cigarettes in past month

http://www.healthindicators.gov/Indicators/Adolescent-use-of-cigarettes-in-past-month---Students-grades-9-through-12_1502/National_0/Profile

Adult Obesity (2010)

Indicator: Percent of Adults (age 18+) who are obese (BMI > 30)

Source: Behavioral Risk Factor Surveillance System 2010

<http://apps.nccd.cdc.gov/BRFSS/list.asp?cat=OB&yr=2010&qkey=4409&state=All>

About this Indicator: Includes all 50 states. National data based on median state prevalence of 50 States and DC (median %=27.6).

Additional Resources and Links:

CDC Vital Signs: Adult Obesity

<http://www.cdc.gov/vitalsigns/AdultObesity/index.html>

MMWR: State-Specific Obesity Prevalence Among Adults --- United States, 2009

http://www.cdc.gov/mmwr/preview/mmwrhtml/mm59e0803a1.htm?s_cid=mm59e0803a1_e%0D%0A

SMART: BRFSS City and County Data (2009): <http://apps.nccd.cdc.gov/BRFSS-SMART/ListMMSAQuest.asp?yr2=2009&MMSA=All&cat=OB&qkey=4409&grp=0>

Health Indicators Warehouse: Obesity (including county level data)

http://www.healthindicators.gov/Indicators/Obesity_15/Profile

Youth Obesity (2009)

Indicator: Percent of high school students who are obese (BMI \geq the 95th percentile for age/sex)

Source: Youth Risk Behavior Surveillance System (YRBSS)

<http://apps.nccd.cdc.gov/YouthOnline/App/Default.aspx?SID=HS>

About this Indicator: National median based on data from 42 states using YBRSS survey of adolescent's grades 9-12 (median%=12) Data not available in CA, DC, IA, MN, NE, OH, OR, VA, and WA.

Additional Resources and Links:

MMWR: Youth Risk Behavior Surveillance—United States, 2009

<http://www.cdc.gov/mmwr/preview/mmwrhtml/ss5905a1.htm>

Adult Seat Belt Use (2009)

Indicator: Percent of observed seat belt use based on probability samples in all 50 states

Source: National Highway Traffic Safety Administration (NHTSA)

<http://www-nrd.nhtsa.dot.gov/Pubs/811324.pdf>



Health Indicator Sortable Stats

About the Data

About this Indicator: NHTSA's National Occupant Protection Use Survey (NOPUS) is a national probability-based survey. National average of 84% includes 50 states, DC, and territories.

Additional Resources and Links:

CDC Vital Signs: Adult Seat Belt Use

<http://www.cdc.gov/vitalsigns/SeatBeltUse/index.html>

MMWR: Nonfatal, Motor Vehicle--Occupant Injuries (2009) and Seat Belt Use (2008) Among Adults --- United States

http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5951a3.htm?s_cid=mm5951a3_w

Youth Seat Belt Use (2009)

Indicator: Percent of high school students who wore a seat belt sometimes, most of the time, or always

Source: Youth Risk Behavior Surveillance System (YRBSS)

<http://apps.nccd.cdc.gov/YouthOnline/App/Default.aspx?SID=HS>

About this Indicator: National median based on data from 40 states using YBRSS survey of adolescent's grades 9-12 (median%=90.3) Data not available in CA, DC, HI, IA, ME, MN, NE, OH, OR, VA, and WA.

Additional Resources and Links:

MMWR: Youth Risk Behavior Surveillance—United States, 2009

<http://www.cdc.gov/mmwr/preview/mmwrhtml/ss5905a1.htm>

Adult Binge Drinking (2010)

Indicator: Percent of Adults (age 18+) who are binge drinkers (males having five or more drinks on one occasion, females having four or more drinks on one occasion)

Source: Behavioral Risk Factor Surveillance System 2010

<http://apps.nccd.cdc.gov/BRFSS/list.asp?cat=AC&yr=2010&qkey=7307&state=All>

About this Indicator: Includes all 50 states and DC. National data based on median state prevalence of 50 States and DC (median %=15.1).

Additional Resources and Links:

CDC Vital Signs: Binge Drinking

<http://www.cdc.gov/vitalsigns/BingeDrinking/index.html>

MMWR: Binge Drinking Among High School Students and Adults --- United States, 2009

http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5939a4.htm?s_cid=mm5939a4_w

SMART: BRFSS City and County Data (2009): <http://apps.nccd.cdc.gov/BRFSS-SMART/ListMMSAQuest.asp?yr2=2009&MMSA=All&cat=AC&qkey=7307&grp=0>

Health Indicators Warehouse: Alcohol: binge drinking (including county level data)

http://www.healthindicators.gov/Indicators/Alcohol-Binge-drinking_17/National_0/Profile

Youth Binge Drinking (2009)

Indicator: High school students reporting having five or more drinks of alcohol in a row within a couple of hours on at least 1 day in last 30 days (adolescents grades 9-12)

Source: Youth Risk Behavior Surveillance System (YRBSS)

<http://apps.nccd.cdc.gov/YouthOnline/App/Default.aspx?SID=HS>



Health Indicator Sortable Stats

About the Data

About this Indicator: National Average based on 41 states using YBRSS survey of adolescent's grades 9-12 (median %=24.2) Data not available in CA, DC, IA, ME, MN, NE, OH, OR, VA, and WA.

Additional Resources and Links:

MMWR: Youth Risk Behavior Surveillance—United States, 2009

<http://www.cdc.gov/mmwr/preview/mmwrhtml/ss5905a1.htm>

Colorectal Cancer Screening (2010)

Indicator: Percent of persons age 50+ who have ever had a sigmoidoscopy or colonoscopy

Source: Behavioral Risk Factor Surveillance System 2010

<http://apps.nccd.cdc.gov/BRFSS/list.asp?cat=CC&yr=2010&qkey=4425&state=All>

About this Indicator: Includes all 50 states and DC. National data based on median state prevalence of 50 States and DC (median %=65.2).

Additional Resources and Links:

CDC Vital Signs: Colorectal Cancer: Cancer Screening

<http://www.cdc.gov/vitalsigns/CancerScreening/ColorectalCancer/index.html>

MMWR: Colorectal Cancer Screening Among Adults Aged 50--75 Years --- United States, 2008

<http://www.cdc.gov/mmwr/preview/mmwrhtml/mm59e0706a1.htm>

Health Indicators Warehouse: Proctoscopy (including county level data)

http://www.healthindicators.gov/Indicators/Proctoscopy_117/Profile

Without Health Care Coverage (2010)

Indicator: Percentage of persons without any kind of health care coverage

Source: Behavioral Risk Factor Surveillance System 2010

<http://apps.nccd.cdc.gov/BRFSS/list.asp?cat=HC&yr=2010&qkey=868&state=All>

About this Indicator: Includes all 50 states and DC. National data based on median state prevalence of 50 States and DC (median %=15).

Additional Resources and Links:

CDC Vital Signs: Access to Health Care

<http://www.cdc.gov/vitalsigns/HealthcareAccess/index.html>

MMWR: Health Insurance Coverage and Health Care Utilization --- United States, 2006--2009 and January--March 2010

http://www.cdc.gov/mmwr/preview/mmwrhtml/mm59e1109a1.htm?s_cid=mm59e1109a1_w

SMART: BRFSS City and County Data (2009): <http://apps.nccd.cdc.gov/BRFSS-SMART/ListMMSAQuest.asp?yr2=2009&MMSA=All&cat=HC&qkey=868&grp=0>

Health Indicators Warehouse: Uninsured (including county level data)

http://www.healthindicators.gov/Indicators/Uninsured_23/Profile

