

Virus Name: Jacareacanga		Abbreviation: JACV
Status Possible Arbovirus	Select Agent No	SALS Level 3
SALS Basis Insufficient experience with virus; i.e., experience factor from SALS surveys was less than 500 in laboratory facilities with low biocontainment.		
Other Information		
Antigenic Group Corriparta		

SECTION I - Full Virus Name and Prototype Number

Prototype Strain Number / Designation BeAr 295042	Accession Number	Original Date Submitted 4/13/1985
Family Orbivirus	Genus	
Information From F.P. Pinheiro and Amelia P.A.T. Rosa	Address Instituto Evandro Chagas, FSESP, Ministry of Health, CP-621, Belem, Para, Brazil	
Information Footnote Reviewed by editor		

Section II - Original Source

Isolated By (name) F. Pinheiro and Amelia P.A.T.Rosa	Isolated at Institute Instituto Evandro Chagas	
Host Genus Cules (Mel) sp., pool of 25 mosquitoes	Species	Host Age/Stage adult
Sex Female		
<u>Isolated From</u>	<u>Isolation Details</u>	
Signs and Symptoms of Illness	Arthropod Engorged	
Time Held Alive before Inoculation 2-3 hours		
Collection Method aspiration from tree trunk	Collection Date 11/11/1975	
Place Collected (Minimum of City, State, Country) Flexal area, km. 212, Itaituba-Jacareacanga		
Latitude 5° 18' S	Longitude 57° 4' W	
Macrohabitat tropical rain forest	Microhabitat ground level	Method of Storage until Inoculated liquid nitrogen and mechanical freezer (-60dC)
Footnotes		

Section III - Method of Isolation

Inoculation Date
5/26/1976

Animal (Details will be in Section 6)
nb mice

Route Inoculated
intracerebral

Reisolation
Not tried

Other Reasons

Homologous Antibody Formation by Source Animal

Test(s) Used

Footnotes

Section IV - Virus Properties

Physicochemical

Pieces (number of genome segments)	Infectivity	Sedimentation Coefficients(s) (S)
Percentage wt, of Virion Protein	Lipid	Carbohydrate
Virion Polypeptides: Number	Details	
Non-virion Polypeptides: Number	Details	
Virion Density	Sedimentation Coefficients(s) (S)	
Nucleocapsid Density	Sedimentation Coefficients(s) (S)	

Stability of Infectivity (effects)

pH (infective range)

Lipid Solvent (ether - % used to test)	After Treatment Titer	Control Titer
Lipid Solvent (chloroform)	After Treatment Titer	Control Titer
Lipid Solvent (deoxycholate) 1:1000	After Treatment Titer 3.3 dex	Control Titer 3.4 dex
Other (formalin, radiation)		

Virion Morphology

Shape	Dimensions	
Mean nm	Range nm	
Measurement Method	Surface Projections/Envelope	Nucleocapsid Dimensions, Symmetry

Morphogenesis

Site of Constituent Formation in Cell	Site of Virion Assembly	Site of Virion Accumulation
Inclusion Bodies	Other	

Hemagglutination

Hemagglutination No	Antigen Source SMB ext. by sucrose-acetone + sonication	Erythrocytes (species used) goose
-------------------------------	---	---

pH Range 5.8 - 7.0	pH Optimum
------------------------------	------------

Temperature Range room, 37dC	Temperature Optimum
--	---------------------

Remarks

Serologic Methods Recommended
CF and NT

Footnotes

No reactions were observed by CF between BeAr 295042 antigen and grouping fluids of A, B, C, Guama, Capim, Bunyamwera, California, Phlebotomus fever, Simbu, Kemerovo, Tacaribe, as well as polyvalents 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, Patois, Bwamba, Congo, Quaranfil, Anopheles A, rabies/LCM etc., and GD VII. It did react with homologous serum, polyvalent Palyam and virus strain BeAr 263191 (isolated in 1974). A relationship has been demonstrated with Corriparta and Acado viruses as follows:

Antigens	Complement-Fixation Test				
	BeAr 263191	BeAr 295042	Sera		
			Acado	Corriparta	Control
BeAr 263191	32/16 ^b	32/16	>=128/256	8/8	0
BeAr 295042	16/16	32/16	>=128/256	8/16	0
Acado	32/16	32/16	>=128/>=256	8/64	0
Corriparta	8/4	16/16	>=128/64	8/16	0
Control	0	0	0	0	0

^b Serum titer/antigen titer; 0 = <4/<4

Neutralization tests performed by the ic route in nb mice showed antigen differences between these agents, as follows:

Virus	Neutralization Test (ic route)			
	BeAr 263191	BeAr 295042	Acado	Corriparta
BeAr 263191	4.4 ^c	3.2	3.3	0.8
BeAr 295042	1.9	3.4	1.3	<=0.2
Acado	2.4	<=1.5	3.7	<=1.3
Corriparta	1.6	<=1.2	<=2.0	2.2

^c LNI in dex

Section VI - Biologic Characteristics

Virus Source (all VERTEBRATE isolates)

Lab Methods of Virus Recovery (ALL ISOLATIONS)
newborn mice

Cell system (a)	Virus passage history (b)	Evidence of Infection							Growth Without CPE +/- (g)
		CPE			PLAQUES				
		Day (c)	Extent (d)	Titer TCD50/ml (e)	Day (c)	Size (f)	Titer PFU/ml (e)		
Vero (CL)	SMB 2		No CPE			No Plaques			

Section VII - Natural Host Range (Additional text can be added below table)

Vertebrate (species and organ) and arthropod	No. isolations/No. tested	No. with antibody/No. tested Test used	Country and region
Culex (Melanoconion) sp.	1/14,110		Km. 212, Itaitube-Jacareacanga road, Para, Brazil
Other Culicidae	0/1,748		
Phlebotomine flies	0/3		

Jacareacanga virus was not isolated from blood or viscera of the following animals captured in this area in 1975: 176 rodents, 63 marsupials, 3 primates, 5 reptiles, 1 ungulate, 1 carnivore and 530 wild birds.

Section VIII - Susceptibility to Experimental Infection (include viremia)

Experimental host and age	Passage history and strain	Inoculation Route-Dose	Evidence of infection	AST (days)	Titer log10/ml
mice (nb)	SMB 5	ic 0.02	death	2.7	
mice (nb)		ip 0.02	none		
mice (nb)		sc			
mice (wn)		ic 0.03	none		
mice (wn)		ip 0.03	none		
mice (nb)	SMB 9	ic 0.02	death		7.2

Section IX - Experimental Arthropod Infection and Transmission

Arthropod species & virus source(a)	Method of Infection log10/ml (b)		Incubation period (c)		Transmission by bite (d)		Assay of arthropod, log10/ml (e)		
	Feeding	Injected	Days	°C	Host	Ratio	Whole	Organ	System

Section X - Histopathology

Character of lesions (specify host)

Inclusion Bodies

Intranuclear

Organs/Tissues Affected

Category of tropism

Section XI - Human Disease

In Nature	Residual	Death
Subclinical	Overt Disease	
Clinical Manifestations		
Number of Cases	Category (i.e. febrile illness, etc.)	

Section XII - Geographic Distribution

Known (Virus detected)
Suspected (Antibody only detected)

Remarks

--