

<b>Virus Name: Jari</b>		<b>Abbreviation: JARIV</b>
Status <b>Possible Arbovirus</b>	Select Agent <b>No</b>	SALS Level <b>3</b>
SALS Basis <b>Insufficient experience with virus; i.e., experience factor from SALS surveys was less than 500 in laboratory facilities with low biocontainment.</b>		
Other Information		
Antigenic Group <b>changuinola</b>		

**SECTION I - Full Virus Name and Prototype Number**

Prototype Strain Number / Designation <b>BeAn 385199</b>	Accession Number	Original Date Submitted <b>4/13/1985</b>
Family <b>orbivirus</b>	Genus	
Information From <b>F.P. Pinheiro and Amelia P.A.T. Rosa</b>	Address <b>Instituto Evandro Chagas, FSESP, Ministry of Health, CP-621, Belem, Para, Brazil</b>	
Information Footnote <b>Reviewed by editor</b>		

**Section II - Original Source**

Isolated By (name) <b>F. Pinheiro and Amelia P.A.T. Rosa (1)</b>	Isolated at Institute <b>Instituto Evandro Chagas</b>	
Host Genus <b>Choloepus didactylus (sloth)</b>	Species	Host Age/Stage <b>adult</b>
Sex <b>Female</b>		
<u>Isolated From</u>	<u>Isolation Details</u>	
<b>Organs/Tissues</b>	<b>pool of heart, liver, spleen, and kidney</b>	
Signs and Symptoms of Illness <b>none</b>	Arthropod	
Time Held Alive before Inoculation		
Collection Method	Collection Date <b>8/4/1980</b>	
Place Collected (Minimum of City, State, Country) <b>Monte Dourado area, Jari, Almerim, Para</b>		
Latitude <b>0° 51' S</b>	Longitude <b>52° 32' W</b>	
Macrohabitat <b>tropical rain forest</b>	Microhabitat <b>canopy</b>	Method of Storage until Inoculated <b>at -60dC</b>
Footnotes		

**Section III - Method of Isolation**

Inoculation Date  
**8/5/1980**

Animal (Details will be in Section 6)  
**nb mice**

Route Inoculated <b>intracerebral</b>	Reisolation <b>Yes</b>
--	---------------------------

Other Reasons

Homologous Antibody Formation by Source Animal

Test(s) Used

Footnotes

**Section IV - Virus Properties**

Physicochemical  
**RNA, Double Strand**

Pieces (number of genome segments) <b>10</b>	Infectivity	Sedimentation Coefficients(s) (S)
Percentage wt, of Virion Protein	Lipid	Carbohydrate
Virion Polypeptides: Number	Details	
Non-virion Polypeptides: Number	Details	
Virion Density	Sedimentation Coefficients(s) (S)	
Nucleocapsid Density	Sedimentation Coefficients(s) (S)	

**Stability of Infectivity (effects)**

pH (infective range)

Lipid Solvent (ether - % used to test)	After Treatment Titer	Control Titer
Lipid Solvent (chloroform)	After Treatment Titer	Control Titer
Lipid Solvent (deoxycholate) <b>1:1000</b>	After Treatment Titer <b>3.8 dex</b>	Control Titer <b>3.6 dex</b>
Other (formalin, radiation)		

**Virion Morphology**

Shape	Dimensions	
Mean nm	Range nm	
Measurement Method	Surface Projections/Envelope	Nucleocapsid Dimensions, Symmetry

**Morphogenesis**

Site of Constituent Formation in Cell

Site of Virion Assembly

Site of Virion Accumulation

Inclusion Bodies

Other

---

**Hemagglutination**

Hemagglutination

Antigen Source

Erythrocytes (species used)

**No**

**SMB ext. by sucrose-acetone + sonication**

**goose**

pH Range

pH Optimum

**5.8 - 7.0**

Temperature Range

Temperature Optimum

**room, 37dC**

Remarks

Serologic Methods Recommended

**CF and NT**

Footnotes

Jari virus was unrelated, by CF, to viruses of groups A, B, C, Bunyamwera and Phlebotomus fever, or to Oropouche, Utinga, Sera do Navio, Melao, Belem, Jurona, Piry, Cocal, Timbo, Chaco, Turlock, Amapari, Flexal, Kwatta, Mosqueiro, Marco, Tembe, Cotia-like, Agua Preta, Ieri, Araguari, Inhangapi, Aruac, Trinita, Pacora, Lukuni, GD VII, Pacui, Acara, Nariva, EMC, mouse hepatitis virus, Jacareacanga, BeAn 306706, Santarem, Para, Cuiaba, Mojui dos Campos, Itupiranga, Mapuera, and herpes simplex viruses. Antigen and antibody (homologous titers, 256/ $\geq$ 128) reacted by CF with Changuinola virus and other viruses of the Changuinola serogroup. Results of neutralization tests with Jari virus and other viruses of the Changuinola serogroup are presented below.

## Neutralization Test (Newborn Mice, ic Route)

Virus	Sera												
	IRI	GUR	OUR	CAN	JAM	ALT	CGL	Purus	Jari	Monte Dourado	Saraca	Almeirim	
Irituia	2.4 <sup>b</sup>						0.6	0.3	0.3	0.0		0.0	0.7
Gurupi		2.6					0.7	$\leq$ 1.7	0.0	0.9		1.0	0.9
Ourem			3.7				0.3	1.2	1.2	0.6		0.7	0.3
Caninde				3.0			$\leq$ 0.3	0.8	0.8	0.1		1.9	0.4
Jamanxi					3.1		$\leq$ 0.6	0.8	$\leq$ 0.2	0.6		$\leq$ 0.2	$\leq$ 0.2
Altamira						3.6	$\leq$ 0.3	1.2	$\leq$ 0.8	1.0		$\leq$ 0.8	$\leq$ 0.5
Changuinola	$\leq$ 1.0	$\leq$ 1.1	$\leq$ 1.0	$\leq$ 1.0	$\leq$ 1.1	$\leq$ 1.0	2.1	$\leq$ 1.0	$\leq$ 1.0	$\leq$ 1.1		$\leq$ 1.0	$\leq$ 1.0
Purus	0.6	0.4	0.6	0.7	0.3	0.4	1.0	2.5	0.0	0.2		$\leq$ 0.5	0.3
Jari	0.6	0.6	0.0	0.4	0.4	0.8	0.3	$\leq$ 0.1	1.7	0.0		0.0	0.0
Monte Dourado	$\leq$ 0.7	$\leq$ 0.7	0.7	1.4	$\leq$ 0.8	$\leq$ 0.7	$\leq$ 0.5	1.2	$\leq$ 0.4	3.0		$\leq$ 0.7	0.8
Saraca	$\leq$ 0.8	$\leq$ 0.8	$\leq$ 0.8	1.7	$\leq$ 0.9	$\leq$ 0.8	$\leq$ 1.2	1.1	$\leq$ 1.0	$\leq$ 0.8		3.0	$\leq$ 1.1
Almeirim	0.0	0.5	$\leq$ 0.3	$\leq$ 0.3	$\leq$ 0.3	$\leq$ 0.3	0.4	0.8	0.0	$\leq$ 0.3		0.2	2.4

<sup>b</sup> LNI in dex

**Section VI - Biologic Characteristics**

Virus Source (all VERTEBRATE isolates)  
heart, liver, spleen, kidney (LV)

Lab Methods of Virus Recovery (ALL ISOLATIONS)  
newborn mice

Cell system (a)	Virus passage history (b)	Evidence of Infection							Growth Without CPE +/- (g)
		CPE			PLAQUES				
		Day (c)	Extent (d)	Titer TCD50/ml (e)	Day (c)	Size (f)	Titer PFU/ml (e)		
Vero (CL)	SMB 3	6	4+				No Plaques		

**Section VII - Natural Host Range (Additional text can be added below table)**

Vertebrate (species and organ) and arthropod	No. isolations/No. tested	No. with antibody/No. tested Test used	Country and region
<i>Choloepus didactylus</i> (organs)	1/21	3/22 NT	Monte Dourado area, Jari, Para, Brazil
<i>Tamandua tetradactyla</i>	0/18	0/7 NT	
<i>Dasypus novemcinctus</i>	0/3	0/4 NT	

Jari virus was not isolated from 6,753 (55 pools) of female *Lutzomyia umbratilis*, 9,665 (79 pools) of male *L. umbratilis*, 516 (10 pools) of female phlebotomines and 1,727 (15 pools) of male phlebotomines collected in the Monte Dourad area in 1980.

**Section VIII - Susceptibility to Experimental Infection (include viremia)**

Experimental host and age	Passage history and strain	Inoculation Route-Dose	Evidence of infection	AST (days)	Titer log <sub>10</sub> /ml
mice (nb)	SMB 4	ic 0.02	illness, death	3.7	
mice (nb)		ip 0.02	survived		
mice (nb)		sc			
mice (wn)		ic 0.03	survived		
mice (wn)		ip 0.03	survived		
mice (nb)	SMB 5	ic 0.02	death		6.1

**Section IX - Experimental Arthropod Infection and Transmission**

Arthropod species & virus source(a)	Method of Infection log <sub>10</sub> /ml (b)		Incubation period (c)		Transmission by bite (d)		Assay of arthropod, log <sub>10</sub> /ml (e)		
	Feeding	Injected	Days	°C	Host	Ratio	Whole	Organ	System

**Section X - Histopathology**

Character of lesions (specify host)

Inclusion Bodies

Intranuclear

Organs/Tissues Affected

Category of tropism

**Section XI - Human Disease**

In Nature	Residual	Death
Subclinical	Overt Disease	
Clinical Manifestations		
Number of Cases	Category (i.e. febrile illness, etc.)	

**Section XII - Geographic Distribution**

Known (Virus detected)
Suspected (Antibody only detected)

**Section XIII - References**

1. Travassos da Rosa, A.P.A., et al. 1984. Intervirology 21:38-49.
--

**Remarks**

--