

| | | |
|--|---------------------------|---------------------------|
| Virus Name: Mermet | | Abbreviation: MERV |
| Status Possible Arbovirus | Select Agent No | SALS Level 2 |
| SALS Basis Results of SALS surveys and information from the Catalogue. | | |
| Other Information | | |
| Antigenic Group Simbu | | |

SECTION I - Full Virus Name and Prototype Number

| | | |
|--|--|---|
| Prototype Strain Number / Designation AV-782 | Accession Number | Original Date Submitted 11/2/1984 |
| Family Bunyaviridae | Genus Bunyavirus | |
| Information From Arbovirology Unit | Address Centers for Disease Control, Atlanta, Georgia 30333, USA | |
| Information Footnote Reviewed by editor | | |

Section II - Original Source

| | | |
|--|--|---|
| Isolated By (name) R.H. Kokernot | Isolated at Institute Urbana, Illinois | |
| Host Genus Progne subis (purple martin) | Species | Host Age/Stage |
| Sex Not Answered | | |
| <u>Isolated From</u> | <u>Isolation Details</u> | |
| Whole Blood | | |
| Signs and Symptoms of Illness None | Arthropod | |
| Time Held Alive before Inoculation | | |
| Collection Method Mist net | Collection Date 7/27/1964 | |
| Place Collected (Minimum of City, State, Country) Metropolis, Massac Co., Illinois | | |
| Latitude 37° 11' N | Longitude 88° 45' W | |
| Macrohabitat Brushy area between village and Ohio River | Microhabitat | Method of Storage until Inoculated Solid CO2, then Revco at -70dC |
| Footnotes | | |

Section III - Method of Isolation

Inoculation Date
8/10/1964

Animal (Details will be in Section 6)
nb mice

| | |
|--|---------------------------------|
| Route Inoculated Intracerebral | Reisolation Not tried |
|--|---------------------------------|

Other Reasons

Homologous Antibody Formation by Source Animal
Not tested

Test(s) Used

Footnotes

Section IV - Virus Properties

Physicochemical
RNA

| | | |
|------------------------------------|--------------------------------------|--------------------------------------|
| Pieces (number of genome segments) | Infectivity | Sedimentation Coefficients(s) (S) |
| Percentage wt, of Virion Protein | Lipid | Carbohydrate |
| Virion Polypeptides: Number | Details | |
| Non-virion Polypeptides: Number | Details | |
| Virion Density | Sedimentation Coefficients(s) (S) | |
| Nucleocapsid Density | Sedimentation Coefficients(s) (S) | |

Stability of Infectivity (effects)

pH (infective range)

| | | |
|--|---|---------------------------------|
| Lipid Solvent (ether - % used to test) | After Treatment Titer | Control Titer |
| Lipid Solvent (chloroform) | After Treatment Titer | Control Titer |
| Lipid Solvent (deoxycholate) 1:200 | After Treatment Titer 2.3 dex | Control Titer 7.0 dex |
| Other (formalin, radiation) | | |

Virion Morphology

| | | |
|--------------------|------------------------------|-----------------------------------|
| Shape | Dimensions | |
| Mean nm | Range nm | |
| Measurement Method | Surface Projections/Envelope | Nucleocapsid Dimensions, Symmetry |

Morphogenesis

| | | |
|---------------------------------------|-------------------------|-----------------------------|
| Site of Constituent Formation in Cell | Site of Virion Assembly | Site of Virion Accumulation |
|---------------------------------------|-------------------------|-----------------------------|

| | |
|------------------|-------|
| Inclusion Bodies | Other |
|------------------|-------|

Hemagglutination

| | | |
|--------------------------------|--|---|
| Hemagglutination Yes | Antigen Source SMB ext. by sucrose-acetone | Erythrocytes (species used) Goose |
|--------------------------------|--|---|

| | |
|----------------------------|--------------------------|
| pH Range 5.8-6.8 | pH Optimum 6.2 |
|----------------------------|--------------------------|

| | |
|--|------------------------------------|
| Temperature Range 4dC - 37dC | Temperature Optimum 37dC |
|--|------------------------------------|

Remarks

HA antigen produced in low titer and obtained only with unlyophilized sucrose-acetone antigen * Determined with BUDR using vaccinia and WEE as controls in Vero cell cultures

Serologic Methods Recommended

CF, NT, HI

Footnotes

HA antigen produced in low titer and obtained only with unlyophilized sucrose-acetone antigen * Determined with BUDR using vaccinia and WEE as controls in Vero cell cultures

HA antigen not inhibited by immune serum or ascitic fluid to the following: arboviruses of groups A, B, C, Guama, Bunyamwera, California, and Turlock as well as Sicilian Sandfly fever, Flanders, Silverwater and Colorado tick fever. In addition, no inhibition of HA was demonstrated with immune ascitic fluids to groups A, C, California, Bunyamwera and Pahayokee-Shark River-Gumbo Limbo (Broad).

CF antigen of AV-782 did not react with immune serum or ascitic fluid to Flanders, yellow fever, St. Louis encephalitis, La Crosse virus, EEE, WEE, Highlands J, Silverwater, Powassan, Tensaw, Colorado tick fever, Turlock, VSNJ, VSI, Cocal, Cowbone Ridge, Tamiami, Anopheles A, Shark River, Pahayokee, trivittatus, Gumbo Limbo, Sawgrass, Newcastle disease virus, encephalomyocarditis, mouse hepatitis, mouse adenovirus, LCM and vaccinia. In addition, immune ascitic fluids to broad groups A, B, C, Guama, Bunyamwera, Capim, Phlebotomus fever, and California failed to react with AV-782 antigen.

IP neutralization tests in suckling mice using another strain of Mermet virus and immune serum to the following did not give evidence of protection: Tensaw, Colorado tick fever, Powassan, Rio Bravo, Modoc, TMA-1381, Kern Canyon, AR-531, CE, EEE, WEE, SLE, Turlock, Flanders, Hart Park and AR-4242.

Cross Reactivities of Mermet (Strain AV-782) Virus with Simbu Group Members (Ht/Ho)

| Registered Virus | Mermet Immune Ascitic Fluid | | | Mermet Antigen | | |
|------------------|-----------------------------|----------|----------|----------------|----------|-------|
| | CF | NT | HI | CF | NT | HI |
| Oropouche | 128/>1024 | /4.3 | | 0/32 | 1.1/3.0 | |
| Sathuperi | 8/>1024 | >1.3/4.3 | | 0/128 | 0.3/>6.0 | |
| Buttonwillow | 32/>1024 | 2.7/4.3 | | 0/64 | 3.5/3.2 | |
| Utinga | 64/>1024 | 1.4/4.3 | | 0/16 | 1.6/2.3 | |
| Akabane | 16/>1024 | 0/4.3 | | 0/128 | 0.6/>4.8 | |
| Ingwavuma | 512/>1024 | >5.5/4.3 | 20/>1280 | 512/32 | 2.8/3.8 | 10/60 |
| Simbu | 8/>1024 | 0.5/4.3 | | 0/16 | 1.3/2.5 | |
| Manzanilla | >1024/>1024 | >3.9/4.3 | 40/>1280 | 16/64 | 3.6/>3.9 | 10/80 |
| Mermet | >1024 | 4.3 | >1280 | >1024 | 4.3 | >1280 |

Mermet virus was shown to be closely related to Ingwavuma and Inini viruses; all three viruses were considered to be subtypes of Manzanilla virus [4].

Section VI - Biologic Characteristics

Virus Source (all VERTEBRATE isolates)
salivary glands (LV)

Lab Methods of Virus Recovery (ALL ISOLATIONS)
newborn mice

| Cell system (a) | Virus passage history (b) | Evidence of Infection | | | | | | | Growth Without CPE +/- (g) |
|-----------------------|------------------------------|-----------------------|---------------|-----------------------|------------|----------|---------------------|--|----------------------------------|
| | | CPE | | | PLAQUES | | | | |
| | | Day (c) | Extent (d) | Titer TCD50/ml (e) | Day (c) | Size (f) | Titer PFU/ml (e) | | |
| BHK-21 (CL) | SMB6 TC4 | | CPE | | 3 | Plaques | 7.6* | | |
| Rabbit kidney (CL) | | | CPE | | 3 | Plaques | 7.3 | | |
| Vero (CL) | | | CPE | | 3 | Plaques | 7.0 | | |
| Duck embryo (PC) | | | CPE | | 3 | Plaques | 5.9 | | |
| * Expressed in dex | | | | | | | | | |

| Vertebrate (species and organ) and arthropod | No. isolations/No. tested | No. with antibody/No. tested Test used | Country and region |
|--|---------------------------|---|--------------------------------|
| Progne subis subis | 1 | | S. Illinois, July 1964 |
| Cyanocitta cristata cristata | 1 | | S. Illinois, Sept. 1964 |
| Hyalocichla ustulata swainsoni | 1 | | S. Illinois, Sept. 1965 |
| Agelaius phoeniceus phoeniceus | 1 | | S. Illinois, May 1964 |
| Richmondia cardinalis (cardinal) | 1/48 | | Dallas, Texas, August 1966 |
| Other wild birds | 0/632 | | Dallas, Texas, Aug.-Sept. 1966 |
| Culex restuans | 1 | | Memphis, Tennessee(3) |
| Culex pipiens | 1 | | |

NT antibody found in wild birds (3,198 tested) collected in Texas, Mississippi, Tennessee, Ohio, Indiana, Illinois and Wisconsin; not in those collected in Kentucky and Missouri (2).

Section VIII - Susceptibility to Experimental Infection (include viremia)

| Experimental host and age | Passage history and strain | Inoculation Route-Dose | Evidence of infection | AST (days) | Titer log ₁₀ /ml | |
|---------------------------|----------------------------|------------------------|-----------------------|------------|-----------------------------|--|
| | | | | | | |
| Mice (nb) | SMB6 TC4 | ic 0.02 | Death | 3 | 7.9 | |
| Mice (nb) | | ip 0.03 | Death | 3-4 | 6.9 | |
| Mice (nb) | | sc | | | | |
| Mice (wn) | | ic 0.02 | Death | 8 | 7.1 | |
| Mice (wn) | | ip 0.03 | Death | 8 | 6.1 | |
| Mice (8 wk) | | ic 0.02 | Antibody | | | |
| Mice (8 wk) | | ip 0.03 | Antibody | | | |

Section IX - Experimental Arthropod Infection and Transmission

| Arthropod species & virus source(a) | Method of Infection log ₁₀ /ml (b) | | Incubation period (c) | | Transmission by bite (d) | | Assay of arthropod, log ₁₀ /ml (e) | | |
|-------------------------------------|---|----------|-----------------------|----|--------------------------|-------|---|-------|--------|
| | Feeding | Injected | Days | °C | Host | Ratio | Whole | Organ | System |
| | | | | | | | | | |
| | | | | | | | | | |

Section X - Histopathology

Character of lesions (specify host)

The only specific findings were in the brain: nonfocal, general increasing glial cells and neuronal degeneration -necrotizing encephalitis. Not a remarkable meningitic or perivascular reaction (suckling mice).

Inclusion Bodies

Intranuclear

Organs/Tissues Affected

Brain (LV)

Category of tropism

Section XI - Human Disease

In Nature

Residual

Death

Subclinical

Overt Disease

Clinical Manifestations

Number of Cases

Category (i.e. febrile illness, etc.)

Section XII - Geographic Distribution

Known (Virus detected)

Colorado; Illinois; Tennessee; Texas,USA

Suspected (Antibody only detected)

Mississippi, Tennessee, Ohio, Indiana, Wisconsin (2)

Section XIII - References

1. Calisher, C.H., et al. 1969. Am. J. Trop. Med. and Hyg. 18:779-788.
2. Calisher, C.H., et al. 1981. Am. J. Trop. Med. and Hyg. 30:473-476.
3. Jakob, W.L., et al. 1971. J. Med. Ent. 16:80-81
4. Kinney, R.M. and Calisher, C.H. 1981. Am. J. Trop. Med. Hyg. 30:1307-1318.

Remarks