

Virus Name: Ngari		Abbreviation: NRIV
Status Probable Arbovirus	Select Agent No	SALS Level
SALS Basis		
Other Information		
Antigenic Group Bunyamwera		

SECTION I - Full Virus Name and Prototype Number

Prototype Strain Number / Designation DAKArD 28542	Accession Number	Original Date Submitted 12/5/1985
Family	Genus Bunyavirus	
Information From J.P. Digoutte and G. Heme	Address Institut Pasteur, BP 220 Dakar, Senegal	
Information Footnote		

Section II - Original Source

Isolated By (name) J.J. Salaun and M. Germain (1)	Isolated at Institute Dakar, Senegal	
Host Genus Aedes simpsoni (males) from eggs collected in nature	Species	Host Age/Stage Adults
Sex Male		
<u>Isolated From</u>	<u>Isolation Details</u>	
Signs and Symptoms of Illness	Arthropod	
Time Held Alive before Inoculation		
Collection Method	Collection Date 1/4/1979	
Place Collected (Minimum of City, State, Country) 10 km from Kedougou, Senegal		
Latitude 12° 36' N	Longitude 12° 15' W	
Macrohabitat Sub-Saharan savannah	Microhabitat Forest gallery	Method of Storage until Inoculated Mechanical freezer at -70dC
Footnotes		

Section III - Method of Isolation

Inoculation Date	
Animal (Details will be in Section 6) nb mice	
Route Inoculated Intracerebral	Reisolation Not tried
Other Reasons	
Homologous Antibody Formation by <u>Source Animal</u>	
Test(s) Used	
Footnotes	

Section IV - Virus Properties

Physicochemical		
Pieces (number of genome segments)	Infectivity	Sedimentation Coefficients(s) (S)
Percentage wt, of Virion Protein	Lipid	Carbohydrate
Virion Polypeptides: Number	Details	
Non-virion Polypeptides: Number	Details	
Virion Density	Sedimentation Coefficients(s) (S)	
Nucleocapsid Density	Sedimentation Coefficients(s) (S)	
<hr/>		
<u>Stability of Infectivity (effects)</u>		
pH (infective range)		
Lipid Solvent (ether - % used to test)	After Treatment Titer	Control Titer
Lipid Solvent (chloroform)	After Treatment Titer 2.8 dex	Control Titer 8.6 dex
Lipid Solvent (deoxycholate)	After Treatment Titer	Control Titer
Other (formalin, radiation)		
<hr/>		
<u>Virion Morphology</u>		
Shape	Dimensions <220 nm	
Mean nm	Range nm	
Measurement Method Millipore filtration	Surface Projections/Envelope	Nucleocapsid Dimensions, Symmetry

Morphogenesis

Site of Constituent Formation in Cell

Site of Virion Assembly

Site of Virion Accumulation

Inclusion Bodies

Other

Hemagglutination

Hemagglutination

Antigen Source

Erythrocytes (species used)

Yes

SMB ext. by sucrose-acetone

Goose

pH Range

pH Optimum

5.0-6.6

6.2

Temperature Range

Temperature Optimum

Room temperature

Remarks

Usual titer = 160

Serologic Methods Recommended

CF, NT

Footnotes

Usual titer = 160

Insitut Pasteur Dakar

CF test-homologous titer = 256/8.

DAKArD 28542 gives positive results with the following immune fluids: Bunyamwera, Ilesha, Shokwe, Biaro, ArD 7343 (Bozo), ArY 357 (Mboke).

Neutralization test eith these viruses gives the following results by the plaque-reduction method on Vero cells.

ANTISERA	VIRUS						
	Ngari	Bunyamwera	Ilesha	Shokwe	Biaro	Bozo(ArB 7343)	Mboke (ArY 357)
Ngari (ArD 28542)	256 *	0	0	0	0	0	0
Bunyamwera	0	256	0	0	0	0	0
Ilesha	0	0	256	0	0	0	0
Shokwe	0	0	0	128	0	0	0
Biaro	0	0	0	0	128	0	0
ArB 7434	0	0	0	0	0	512	0
ArY 357	0	0	0	0	0	0	256

* Reciprocal of highest antiserum dilution producing > 90% plaque inhibition; 0 = <16.

Results indicate that DAKArD 28542 is apparently a new virus.

Section VI - Biologic Characteristics

Virus Source (all VERTEBRATE isolates)

Lab Methods of Virus Recovery (ALL ISOLATIONS)
Newborn mice

Cell system (a)	Virus passage history (b)	Evidence of Infection						
		CPE			PLAQUES			Growth Without CPE +/- (g)
		Day (c)	Extent (d)	Titer TCD50/ml (e)	Day (c)	Size (f)	Titer PFU/ml (e)	
Vero (CL)	SM 4	2-3	CPE		3	2	6.4 dex	

Section VII - Natural Host Range (Additional text can be added below table)

Vertebrate (species and organ) and arthropod	No. isolations/No. tested	No. with antibody/No. tested Test used	Country and region
Aedes simpsoni (males)	1		Kendougou, Senegal
Aedes vittatus	2		Kendougou, Senegal
Aedes neoafricanus	1		Kendougou, Senegal
Aedes Argenteopunctatus	1		Kendougou, Senegal
Anopheles gambiae	5		Burkino Faso (2)
Anopheles gamibiae	1		Bozo, Central African Republic (2)
Anopheles mascarensis	1		Madagascar (2)

Section VIII - Susceptibility to Experimental Infection (include viremia)

Experimental host and age	Passage history and strain	Inoculation Route-Dose	Evidence of infection	AST (days)	Titer log ₁₀ /ml
mice (nb)	SM 4	ic	Death	2	10.3
"" (nb)		ip	Death	3	
"" (nb)		sc			
"" (wn)		ic	Death	6	
"" (wn)		ip	Antibodies		

Section IX - Experimental Arthropod Infection and Transmission

Arthropod species & virus source(a)	Method of Infection log ₁₀ /ml (b)		Incubation period (c)		Transmission by bite (d)		Assay of arthropod, log ₁₀ /ml (e)		
	Feeding	Injected	Days	°C	Host	Ratio	Whole	Organ	System

Section X - Histopathology

Character of lesions (specify host)

Inclusion Bodies

Intranuclear

Organs/Tissues Affected

Category of tropism

Section XI - Human Disease

In Nature	Residual	Death
Subclinical	Overt Disease	
Clinical Manifestations		
Number of Cases	Category (i.e. febrile illness, etc.)	

Section XII - Geographic Distribution

Known (Virus detected) Senegal, Burkino Faso, Central African Republic, Madagascar
Suspected (Antibody only detected)

Section XIII - References

1. Institut Pasteur Dakar Annual Report. 1980. 2. Institut Pasteur Dakar Annual Report. 1984.
--

Remarks

--