

<b>Virus Name: Cacipacore</b>		<b>Abbreviation: CPCV</b>
Status <b>Possible Arbovirus</b>	Select Agent <b>No</b>	SALS Level <b>3</b>
SALS Basis <b>Insufficient experience with virus; i.e., experience factor from SALS surveys was less than 500 in laboratory facilities with low biocontainment.</b>		
Other Information		
Antigenic Group <b>B</b>		

#### SECTION I - Full Virus Name and Prototype Number

Prototype Strain Number / Designation <b>BeAn 327600</b>	Accession Number	Original Date Submitted <b>9/24/1984</b>
Family <b>Flaviviridae</b>	Genus <b>Flavivirus</b>	
Information From <b>F.P. Pinheiro and Amelia P.A.T. Rosa</b>	Address <b>Instituto Evandro Chagas, FSESP, Ministry of Health, CP-621, Belem,Para,Brazil</b>	
Information Footnote		

#### Section II - Original Source

Isolated By (name) <b>F.P. Pinheiro and Amelia P.A.T. Rosa</b>	Isolated at Institute <b>Instituto Evandro Chagas</b>	
Host Genus <b>Formicarius analis (black-faced ant bird)</b>	Species	Host Age/Stage <b>Adult</b>
Sex <b>Male</b>		
<u>Isolated From</u>  <b>Whole Blood</b>	<u>Isolation Details</u>	
Signs and Symptoms of Illness	Arthropod	
Time Held Alive before Inoculation		
Collection Method <b>Netted</b>	Collection Date <b>7/28/1997</b>	
Place Collected (Minimum of City, State, Country) <b>Cachoeira Porteira, km.71, Oriximina, Para, Brazil</b>		
Latitude <b>0° 35' S</b>	Longitude <b>56° 45' W</b>	
Macrohabitat <b>Tropical Rain Forest</b>	Microhabitat <b>Ground Level</b>	Method of Storage until Inoculated <b>Liquid nitrogen and mechanical freezer (-60C)</b>
Footnotes		

### Section III - Method of Isolation

Inoculation Date

**11/18/1977**

Animal (Details will be in Section 6)

**nb mice**

Route Inoculated

**intracerebral**

Reisolation

**No**

Other Reasons

Homologous Antibody Formation by Source Animal

Test(s) Used

Footnotes

### Section IV - Virus Properties

Physicochemical

Pieces (number of genome segments)	Infectivity	Sedimentation Coefficients(s) (S)
Percentage wt, of Virion Protein	Lipid	Carbohydrate
Virion Polypeptides: Number	Details	
Non-virion Polypeptides: Number	Details	
Virion Density	Sedimentation Coefficients(s) (S)	
Nucleocapsid Density	Sedimentation Coefficients(s) (S)	

#### Stability of Infectivity (effects)

pH (infective range)

Lipid Solvent (ether - % used to test)	After Treatment Titer	Control Titer
Lipid Solvent (chloroform)	After Treatment Titer	Control Titer
Lipid Solvent (deoxycholate) <b>1:1000</b>	After Treatment Titer <b>&lt;=0.5 dex</b>	Control Titer <b>4.0 dex</b>
Other (formalin, radiation)		

#### Virion Morphology

Shape	Dimensions	
Mean nm	Range nm	
Measurement Method	Surface Projections/Envelope	Nucleocapsid Dimensions, Symmetry

**Morphogenesis**

Site of Constituent Formation in Cell	Site of Virion Assembly	Site of Virion Accumulation
Inclusion Bodies	Other	

**Hemagglutination**

Hemagglutination <b>Yes</b>	Antigen Source <b>SMB ext. by sucrose-acetone</b>	Erythrocytes (species used) <b>Goose</b>
pH Range <b>5.8-7.0</b>	pH Optimum <b>7.0</b>	
Temperature Range <b>Room, 37C</b>	Temperature Optimum	
Remarks		
Serologic Methods Recommended <b>CF, HI, and NT</b>		
Footnotes		

Serological studies conducted in Brazil demonstrated that Cacipacore virus was a group B arbovirus, and that it was different from other known flaviviruses of the Amazon (see tables below). At YARU, additional CF tests indicated it was in the JE-MVE-WN complex, but different from any of the known members tested [1]. Subsequent additional cross-neutralization tests with 63 other flaviviruses showed that Cacipacore virus was distinct from all other flaviviruses. It did not share a close antigenic relationship with any of the flaviviruses, and thus could not be placed in any of the complexes comprising serogroup B [2].

Antigens	Complement-Fixation Test						Group B
	BeAn 327600	YF	BSQ	ILH	SLE	ROC	
BeAn 327600	16/>64 <sup>a</sup>	0	0	4/4	4/16	16/4	16/4
Yellow fever	0	64/>64					16/>64
Bussuquara	0		64/>64				32/>64
Ilheus	0			64/>64			64/>64
SLE	4/64				64/>64		>128/64
Rocio	0					>128/64	16/>64

<sup>a</sup> Serum titer/antigen titer: 0 - <4/<4

Viruses	Neutralization Test (Newborn Mice, ic Route)						
	BeAn 327600	YF	BSQ	ILH	SLE	ROC	Group B
BeAN 327600	4.7 <sup>b</sup>	1.0	0.9	2.1	1.9	2.3	2.8
Yellow fever		4.2					
Bussuquara	0.5		3.5				3.4
Ilheus	<=1.0			5.0			3.7
SLE	0.7				3.0		3.0
Rocio	<=1.9					5.0	<=3.9

<sup>b</sup> LNI in dex

## Section VI - Biologic Characteristics

Virus Source (all VERTEBRATE isolates)  
Blood (LV)

Lab Methods of Virus Recovery (ALL ISOLATIONS)  
Newborn mice

Cell system (a)	Virus passage history (b)	Evidence of Infection									
		CPE			PLAQUES			Growth Without CPE +/- (g)			
		Day (c)	Extent (d)	Titer TCD50/ml (e)	Day (c)	Size (f)	Titer PFU/ml (e)				
Vero (CL)	P-unk. SM 1				8	Plaques	7.2 °(2)				
LLC-MK2 (CL)						No plaques	(2)				
Duck embryo (PC)						No plaques	(2)				
° Expressed in dex											

## Section VII - Natural Host Range (Additional text can be added below table)

Vertebrate (species and organ) and arthropod	No. isolations/No. tested	No. with antibody/No. tested Test used	Country and region
Wild birds (blood)	1/2,812	14/1,310 HI	Cachoeira Porteira area, Para, Brazil, 1976-79
Rodents	0/377	1/156 HI	
Marsupials	0/215	0/78 HI	
Primates	0/206	0/133 HI	
Bats	0/260	0/45 HI	
Carivores	0/13	0/3 HI	
Ungulates	0/37	0/8 HI	
Edentates	0/7	0/4 HI	
Reptiles	0/47	0/20 HI	



Reptiles	0/47	0/20 HI	
Birds		8/790 HI	Porto Trometas, Para, Brazil, 1979-80
Marsupials		0/40 HI	
Rodents		0/75 HI	
Primates		0/34 HI	
Bats		1/95 HI	
Reptiles		0/48 HI	
Edentates		0/7 HI	
Ungulates and carnivores		0/8 HI	
Bats		0/50 HI	Santa Isabel do Para, Brazil, 1979
Bats		0/43 HI	Uruacu and Barro Alto, Para, Brazil, 1980
Rodents, marsupials and primates		0/9 HI	
Birds		2/428 HI	Tome Acu, Para, Brazil, 1978
Birds		1/143 HI	Santa Isabel do Para, Brazil, 1979
Man		2/2,572 HI	Para (several areas), 1977-80
Man		0/246 HI	Amazonas (several areas), 1979
Man		0/622 HI	Goiias (several areas), 1980
Man (Xicrin Indians)		0/49 HI	Para, 1978
<p>Cacipacore virus was not isolated from the following arthropods captured in the Cachocira Porteira area, from 1976-79: 382 Sabethes (39 pools), 14,334 other Culicidae (2,252 pools) and 3,015 phlebotomines (58 pools).</p>			

## Section VIII - Susceptibility to Experimental Infection (include viremia)

Experimental host and age	Passage history and strain	Inoculation Route-Dose	Evidence of infection	AST (days)	Titer log10/ml
mice (nb)	SMB 4	ic 0.02	illness, death	4.0	
"" (nb)		ip 0.02	illness, death	9.8	
"" (nb)		sc			
"" (wn)		ic 0.03	illness, death	6.0	
"" (wn)		ip 0.03	antibody		
"" (nb)	SMB 8	ic 0.02	death		9.1

## Section IX - Experimental Arthropod Infection and Transmission

Arthropod species & virus source(a)	Method of Infection log10/ml (b)		Incubation period (c)		Transmission by bite (d)		Assay of arthropod, log10/ml (e)		
	Feeding	Injected	Days	°C	Host	Ratio	Whole	Organ	System

## Section X - Histopathology

Character of lesions (specify host)

Inclusion Bodies

Intranuclear

Organs/Tissues Affected

Category of tropism

**Section XI - Human Disease**

In Nature	Residual	Death
Subclinical	Overt Disease	
Clinical Manifestations		
Number of Cases	Category (i.e. febrile illness, etc.)	

**Section XII - Geographic Distribution**

Known (Virus detected)  
**Para**

Suspected (Antibody only detected)

**Section XIII - References**

1. Shope, R.E. Personal communication. 1981.  
2. Calisher, C.H. Personal communication. 1982.

**Remarks**